

Principal Component Analysis

Question 1: PCA by hand

Consider a data matrix given by

$$\mathbf{X} = \begin{pmatrix} 24 & 22 & 24 \\ 24 & 21 & 25 \\ 24 & 22 & 20 \\ 24 & 23 & 21 \end{pmatrix}.$$

- Derive the principal components via eigen decomposition of the sample covariance matrix.
- Let us assume that we want to reduce the data's dimension to $k = 2$. Calculate the new data points in \mathbb{R}^2 .

Solution:

1. **Compute the sample covariance matrix:**

Recall from the lecture, that the following holds for the sample covariance matrix:

$$\mathbf{S} = \frac{1}{n-1} \mathbf{X}_C^\top \mathbf{X}_C = \frac{1}{n-1} \sum_{i=1}^n (\mathbf{x}_i - \bar{\mathbf{x}}) (\mathbf{x}_i - \bar{\mathbf{x}})^\top.$$

In this case, we have

$$\bar{\mathbf{x}} = \begin{pmatrix} \frac{1}{4} (24 + 24 + 24 + 24) \\ \frac{1}{4} (22 + 21 + 22 + 23) \\ \frac{1}{4} (24 + 25 + 20 + 21) \end{pmatrix} = (24, 22, 22.5)^\top.$$

It follows that

$$\begin{aligned}
\mathbf{S} &= \frac{1}{3} \left(\begin{pmatrix} (24-24) \\ (22-22) \\ (24-22.5) \end{pmatrix} \left((24-24), (22-22), (24-22.5) \right) + \right. \\
&\quad \begin{pmatrix} (24-24) \\ (21-22) \\ (25-22.5) \end{pmatrix} \left((24-24), (21-22), (25-22.5) \right) + \\
&\quad \begin{pmatrix} (24-24) \\ (22-22) \\ (20-22.5) \end{pmatrix} \left((24-24), (22-22), (20-22.5) \right) + \\
&\quad \left. \begin{pmatrix} (24-24) \\ (23-22) \\ (21-22.5) \end{pmatrix} \left((24-24), (23-22), (21-22.5) \right) \right) \\
&= \frac{1}{3} \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 2.25 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -2.50 \\ 0 & -2.5 & 6.25 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 6.25 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1.50 \\ 0 & -1.5 & 2.25 \end{pmatrix} \right) \\
&= \frac{1}{3} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2 & -4 \\ 0 & -4 & 17 \end{pmatrix}.
\end{aligned}$$

2. Perform an eigen decomposition of \mathbf{S} :

First, we need to compute the eigenvalues via the characteristic polynomial

$$\det(\mathbf{S} - \lambda \mathbf{I}_3) \stackrel{!}{=} 0.$$

$$\begin{aligned}
\Rightarrow \begin{vmatrix} -\lambda & 0 & 0 \\ 0 & \left(\frac{2}{3} - \lambda\right) & -\frac{4}{3} \\ 0 & -\frac{4}{3} & \left(\frac{17}{3} - \lambda\right) \end{vmatrix} &= (-\lambda) \left(-\lambda + \frac{2}{3}\right) \left(-\lambda + \frac{17}{3}\right) - \left(-\lambda \cdot -\frac{4}{3} \cdot -\frac{4}{3}\right) \\
&= -\lambda^3 + \frac{19}{3}\lambda^2 - \frac{34}{9}\lambda - \left(-\frac{16}{9}\lambda\right) = \\
&= -\lambda^3 + \frac{19}{3}\lambda^2 - 2\lambda \stackrel{!}{=} 0.
\end{aligned}$$

$$\Rightarrow \lambda_1 = 6, \quad \lambda_2 = \frac{1}{3}, \quad \lambda_3 = 0.$$

Eigenvector corresponding to λ_1

$$\begin{pmatrix} -6x_1 & 0 & 0 \\ 0 & -\frac{16}{3}x_2 & -\frac{4}{3}x_3 \\ 0 & -\frac{4}{3}x_2 & -\frac{1}{3}x_3 \end{pmatrix} \stackrel{!}{=} 0 \quad \Leftrightarrow \quad v_1 = \begin{pmatrix} 0 \\ -\frac{1}{4} \\ 1 \end{pmatrix}.$$

v_1 needs to be normalized: $v_1 = (0.0000000, -0.2425356, 0.9701425)^\top$

Eigenvector corresponding to λ_2

$$\begin{pmatrix} -\frac{1}{3}x_1 & 0 & 0 \\ 0 & -\frac{1}{3}x_2 & -\frac{4}{3}x_3 \\ 0 & -\frac{4}{3}x_2 & -\frac{16}{3}x_3 \end{pmatrix} \stackrel{!}{=} 0 \Leftrightarrow v_2 = \begin{pmatrix} 0 \\ 4 \\ 1 \end{pmatrix}.$$

v_2 needs to be normalized: $\mathbf{v}_2 = (0.0000000, 0.9701425, 0.2425356)^\top$

Eigenvector corresponding to λ_3

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{2}{3}x_2 & -\frac{4}{3}x_3 \\ 0 & -\frac{4}{3}x_2 & \frac{17}{3}x_3 \end{pmatrix} \stackrel{!}{=} 0 \Leftrightarrow v_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

v_3 does not need to be normalized.

Finally, the eigen decomposition of \mathbf{S} is given by

$$\mathbf{S} = (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) \begin{pmatrix} 6 & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \mathbf{v}_1^\top \\ \mathbf{v}_2^\top \\ \mathbf{v}_3^\top \end{pmatrix}$$

and the PCs are $\mathbf{a}_1 = \mathbf{v}_1$, $\mathbf{a}_2 = \mathbf{v}_2$, and $\mathbf{a}_3 = \mathbf{v}_3$.

- b) To use the PCs for dimension reduction, we multiply the original data with the matrix of the first k columns of eigenvectors.

In our case, $k = 2$, so we achieve dimension reduction via

$$\mathbf{X}(\mathbf{v}_1, \mathbf{v}_2) = \begin{pmatrix} 24 & 22 & 24 \\ 24 & 21 & 25 \\ 24 & 22 & 20 \\ 24 & 23 & 21 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ -0.2425356 & 0.9701425 \\ 0.9701425 & 0.2425356 \end{pmatrix} = \begin{pmatrix} 17.94764 & 27.16399 \\ 19.16031 & 26.43638 \\ 14.06707 & 26.19385 \\ 14.79467 & 27.40653 \end{pmatrix}.$$

Question 2: Invariance of PCA w.r.t. transform

Given a PCA of a data matrix $\mathbf{X} \in \mathbb{R}^{n \times m}$, consider the matrix of scores

$$\mathbf{Y} = \begin{pmatrix} y_{11} & \dots & \dots & y_{n1} \\ \vdots & \vdots & \vdots & \vdots \\ y_{1m} & \dots & \dots & y_{nm} \end{pmatrix} = [\mathbf{y}_1, \dots, \mathbf{y}_n]^\top \in \mathbb{R}^{m \times n},$$

where each columns gives the coordinates \mathbf{y}_i of observation i , $i = 1, \dots, n$, in the m -dimensional space with the principal component (vectors) as axes.

- a) Show that the sample covariance of \mathbf{Y} is equal to $\mathbf{\Lambda}_{\text{ord}}$, i.e. the diagonal matrix of ordered eigenvalues of either the sample covariance matrix \mathbf{S} .
- b) In the lecture, we have learned that PCA is not scale-invariant when we solve the optimization problem $\mathbf{a}_p^\top \mathbf{S} \mathbf{a}_p \rightarrow \max$, only when we solve $\mathbf{a}_p^\top \mathbf{R} \mathbf{a}_p \rightarrow \max$.

Can you reason why this is the case, using a diagonal matrix $\mathbf{T} \in \mathbb{R}^{m \times m}$ which transforms the variable scales by replacing each observation \mathbf{x}_i with $\mathbf{T}\mathbf{x}_i$?

- c) Next, consider shifting each data point by a constant $c \in \mathbb{R}$. Is PCA invariant w.r.t. a shift of each data point by a constant?
- d) Lastly, consider an orthogonal matrix $\mathbf{A} \in \mathbb{R}^{m \times m}$. How does PCA behave w.r.t. orthogonal transformation, i.e. w.r.t. replacement of each observation \mathbf{x}_i with $\mathbf{A}\mathbf{x}_i$?

Solution:

a) Let $\mathbf{S} \in \mathbb{R}^{m \times m}$ again denote the sample covariance matrix for the following.

We recall that

- For \mathbf{V} denoting the matrix whose columns are the eigenvectors of \mathbf{S} , ordered in descending order according to the corresponding eigenvalues and \mathbf{X}_C denoting the centered data matrix, we have
 - $\mathbf{S} = \mathbf{V}\mathbf{\Lambda}_{\text{ord}}\mathbf{V}^\top$ and
 - $\mathbf{X}_C = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^\top$.
- For \mathbf{a}_p denoting the p th PC (i.e. p th column of \mathbf{V}), $p = 1, \dots, m$, the p th entry of \mathbf{y}_i is given by

$$y_{ip} = \mathbf{a}_p^\top (\mathbf{x}_i - \bar{\mathbf{x}}), \quad i = 1, \dots, n$$

$$\Leftrightarrow \mathbf{y}_i = (y_{i1}, y_{i2}, \dots, y_{im})^\top = \mathbf{V}^\top (\mathbf{x}_i - \bar{\mathbf{x}}), \quad i = 1, \dots, n.$$

It immediately follows that the sample covariance matrix of \mathbf{Y} is given by

$$\begin{aligned} \frac{1}{n-1} \sum_{i=1}^n \mathbf{y}_i \mathbf{y}_i^\top &= \frac{1}{n-1} \sum \mathbf{V}^\top (\mathbf{x}_i - \bar{\mathbf{x}}) (\mathbf{x}_i - \bar{\mathbf{x}})^\top \mathbf{V} \\ &= \mathbf{V}^\top \mathbf{S} \mathbf{V} \\ &= \mathbf{V}^\top \mathbf{V} \mathbf{\Lambda}_{\text{ord}} \mathbf{V}^\top \mathbf{V} \text{ substituting the eigen decomposition for } \mathbf{S} \\ &= \mathbf{\Lambda}_{\text{ord}}. \end{aligned}$$

b) Just as in the first exercise, that the following holds for the sample covariance matrix:

$$\mathbf{S} = \frac{1}{n-1} \mathbf{X}_C^\top \mathbf{X}_C = \frac{1}{n-1} \sum_{i=1}^n (\mathbf{x}_i - \bar{\mathbf{x}}) (\mathbf{x}_i - \bar{\mathbf{x}})^\top.$$

Now, if we change the scale of a data matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ by replacing each observation \mathbf{x}_i with $\mathbf{T}\mathbf{x}_i$, the new data's arithmetic mean is given by $\frac{1}{n} \sum_{i=1}^n \mathbf{T}\mathbf{x}_i = \mathbf{T}\bar{\mathbf{x}}$ and the new data's sample covariance matrix by

$$\begin{aligned} &\frac{1}{n-1} \sum_{i=1}^n (\mathbf{T}\mathbf{x}_i - \mathbf{T}\bar{\mathbf{x}}) (\mathbf{T}\mathbf{x}_i - \mathbf{T}\bar{\mathbf{x}})^\top \\ &= \frac{1}{n-1} \sum_{i=1}^n \mathbf{T} (\mathbf{x}_i - \bar{\mathbf{x}}) (\mathbf{x}_i - \bar{\mathbf{x}})^\top \mathbf{T}^\top \\ &= \mathbf{T} \mathbf{S} \mathbf{T}^\top \quad \text{because } \mathbf{T} \text{ is diagonal} \quad \mathbf{T} \mathbf{S} \mathbf{T}^\top. \end{aligned}$$

Clearly, the eigenvalues and eigenvectors of \mathbf{TST} will not be identical to those of \mathbf{S} unless all diagonal entries are equal to 1.

\Rightarrow This is what is meant by "PCA is not scale-invariant".

However, since the sample correlation has standardized entries, this is not an issue when considering \mathbf{R} instead of \mathbf{S} .

c) PCA is invariant w.r.t. shifting by a constant, even when using the sample covariance matrix \mathbf{S} .

This holds because, each point of the shifted data is given by $\mathbf{x}_i + c$, the arithmetic mean by $\bar{\mathbf{x}} + c$, and the shifted data's sample covariance matrix by

$$\begin{aligned} & \frac{1}{n-1} \sum_{i=1}^n \left((\mathbf{x}_i + c) - (\bar{\mathbf{x}} + c) \right) \left((\mathbf{x}_i + c) - (\bar{\mathbf{x}} + c) \right)^\top \\ &= \frac{1}{n-1} \sum_{i=1}^n (\mathbf{x}_i - \bar{\mathbf{x}}) (\mathbf{x}_i - \bar{\mathbf{x}})^\top \\ &= \mathbf{S}. \end{aligned}$$

Clearly, it immediately follows that we get the same score vectors.

This is also easily shown:

$$\mathbf{y}_i = \mathbf{V}^\top (\mathbf{x}_i + \mathbf{c} - (\bar{\mathbf{x}} + \mathbf{c})) = \mathbf{V}^\top (\mathbf{x}_i - \bar{\mathbf{x}}) \quad \forall i \in \{1, \dots, n\}.$$

d) Equivalently to subtask a), the sample covariance matrix of the orthogonally transformed data, i.e. data with new observations \mathbf{Ax}_i , is given by

$$\begin{aligned} & \frac{1}{n-1} \sum_{i=1}^n (\mathbf{Ax}_i - \mathbf{A}\bar{\mathbf{x}}) (\mathbf{Ax}_i - \mathbf{A}\bar{\mathbf{x}})^\top \\ &= \frac{1}{n-1} \sum_{i=1}^n \mathbf{A} (\mathbf{x}_i - \bar{\mathbf{x}}) (\mathbf{x}_i - \bar{\mathbf{x}})^\top \mathbf{A}^\top \\ &= \mathbf{ASA}^\top. \end{aligned}$$

Thereby, PCA is definitely not invariant w.r.t. orthogonal transformation.

However, the following also holds for the above sample covariance matrix:

$$\mathbf{ASA}^\top = \mathbf{AV}\mathbf{\Lambda}\mathbf{V}^\top \mathbf{A}^\top = \mathbf{B}\mathbf{\Lambda}\mathbf{B}^\top,$$

for $\mathbf{B} := \mathbf{AV}$. Since \mathbf{B} is also orthogonal by definition, it follows that the eigenvalues of the sample covariance matrix **are not changed by the orthogonal transformation!**

For this reason, PCA is sometimes called *equivariant with respect to orthogonal transformations*.

Question 3: Interpreting PCA output in R

There are two main ways to perform PCA in R:

- the `princomp()` function - based on eigen decomposition and
- the `prcomp()` function - based on singular value decomposition (SVD).

According to the R help, `prcomp()` via SVD has slightly better numerical accuracy. Here you can use the option `scale=TRUE` to perform standardized PCA, i.e. the version that iteratively solves $\mathbf{a}_p^\top \mathbf{R} \mathbf{a}_p \rightarrow \max$.

For visualization of PCA results, the `factoextra` package is very popular; except for biplots, for which the `ggfortify` package is standard.

- a) Perform PCA on the `iris` data set excluding the variable `Species` and interpret the output.
- b) Plot the scree plot and select the number of PCs that should be selected for dimension reduction according to each of the criteria on lecture-slide 67.
- c) Plot the Biplot and interpret it.

Solution:

a) *See R code.*

The output is

Standard deviations (1, ..., p=4):

```
[1] 1.7083611 0.9560494 0.3830886 0.1439265
```


Rotation (n x k) = (4 x 4):

	PC1	PC2	PC3	PC4
Sepal.Length	0.5210659	-0.37741762	0.7195664	0.2612863
Sepal.Width	-0.2693474	-0.92329566	-0.2443818	-0.1235096
Petal.Length	0.5804131	-0.02449161	-0.1421264	-0.8014492
Petal.Width	0.5648565	-0.06694199	-0.6342727	0.5235971

Interpretation:

- The standard deviations are the standard deviations of the principal components, which are equal to the square roots of the eigenvalues of the covariance/correlation matrix.
- The Rotation columns are equal to the principal component (vectors).
Meanwhile, the Rotation rows correspond to the loadings of each variable.

b) *See R code.*



- *Kaiser criterion*: Principal components with eigenvalue greater than 1. (I.e. the maximal k s.t. $\lambda_k > 1$) – **The choice would be 1.**
- All principal components needed to get a total of 80% of the variance. (I.e. the minimal k s.t. $\text{tr}(\mathbf{\Lambda}_{\text{ord}})^{-1} \cdot \sum_{i=1}^k \lambda_k \geq 0.8$) – **The choice would again be 2.**
- *Scree Plot*: Consider a graphical representation of the eigenvalues. Use as many principal components up to the bend of the graph (elbow).
 - **Here, we might decide to go with 3.**
- Simply choose k so that it is convenient (e.g. for a planned visualization).
 - **For visualization, one would often choose 2.**

c) See R code.

The Biplot overlays scoreplots, i.e. dots with coordinates given by the first k entries of the score vectors \mathbf{y}_i , $i = 1, \dots, n$, with loading plots, i.e. arrows that point towards the coordinates given by the first k entries of the columns of the Rotation matrix from subtask a).

In the plot below, we can observe a few things:

- The data may vaguely be divided into two clusters on the first component.
- Since the loadings for `Petal.Length` and `Petal.Width` mostly contribute to the variability along PC1.
- `Petal.Length` and `Petal.Width` are highly positively correlated with each other, but both negatively correlated with `Sepal.Width`.

